Seizure Emergency Care Plan and Medication PARENT/GUARDIAN complete and sign the t		1
	th date:	+
	one:	Dlaga
	one:	Place child's
3 - 1 - 1 - 1 - 1 - 1 - 1 - 1 - 1 - 1 -	ade:	photo here
Triggers: ☐ tiredness ☐ flashing lights ☐ illness		_ prioto riere
Seizure Aura (if any): Seizure history: Convulsive Focal Ab		
Describe:	Selice Date of last kilowit selzure	
Antiseizure Medication Taken at Home Common si	ide effects	
Other Seizure Treatments/Special Diet Therapy:		
give permission for school personnel to share this informatio necessary, contact our health care provider. I assume full resp approve this Seizure Emergency Care Plan for my child.	n, follow this plan, administer medication and care for my chi onsibility for providing the school with prescribed medication	Id and, if n and devices. I 504 plan
PARENT SIGNATURE DATE	SCHOOL NURSE SIGNATURE DATE	_ □ IEP
HEALTH CARE PROVIDER to complete all items, S IF YOU SEE THIS:	SIGN and DATE completed form. DO THIS:	
Convulsive Generalized Tonic Clonic:	Time the seizure	
You will see loss of consciousness. Stiffening of the body. Rhythmic jerking movements. Convulsive seizures may last 1-5 minutes. The child may have a warning (aura) before the seizure. Sleepiness and confusion may occur after the seizure.	 Keep calm. Provide reassurance. Protect head, keep airway clear, turn on side if possible. Do not place anything in mouth. Call 911 if student is injured or has difficulty breathing. Call parent. Stay with student until recovered from seizure. Administer rescue treatments as marked below. 	
These seizures may begin with an aura. They may be partly alert or unconscious. You may see lip smacking, chewing, eye blinking, or picking at clothes. These seizures usually last 1-2 minutes.	 Time the seizure Gently guide child away from danger. Stay with student and reassure them until recovered from seizure. Do not treat staring that is stopped by a touch or a nudge. Call parent. Administer rescue treatments as marked below. 	
Absence: You will see quick changes in alertness. May see eye flutter or small twitching. Usually last less than 10 seconds.		
Rescue Treatments		
☐ Child has a VNS. Child/staff may swipe with aura. Staff Give rescue medications below if seizure does no		seizure stops.
If seizure <u>lasts longer</u> than minutes administer:		
☐ Diastatmg rectally ☐ Midaz	olammg in the nose	cheek
☐ Multistep seizure rescue plan – Please see att	ached letter for details.	
If <u>cluster</u> of or more seizures in min administer:		
☐ Diastatmg rectally ☐ Midaz	olammg in the nose	cheek
☐ Multistep seizure rescue plan – Please see att	ached letter for details.	
If emergency medication is administered: Call 911 in the control of the control	immediately or ☐ Call 911 if seizure does not stop witl	nin 5 minutes
	shild is aversionaine estimate	
If no emergency medication is at school and the Call family to bring medications to school or pick up to	child. Call EMS if seizure lasts more than min	
Accommodations: Always take seizure action plan a Close adult supervision when swimming or climbing.		s and field trips.
HEALTH CARE PROVIDER SIGNATURE PRINT	ROVIDER'S NAME PHONE/FAX I	DATE